


**Gregorio Romero-González^{a,b}, Fredzzia Graterol^a, Jordi Soler^a, Ines Perezpayá^a, Maribel Troya^a, Nestor Rodriguez-Chitiva^a, Rosa Lopez^c, Ana Cia-Hidalgo^c, Marina Urrutia^a, Javier Paul-Martínez^a, Anna Vila^a, Jordi Bover^a, Jordi Ara^a (on behalf of the BIOSONO Marker study group)*

a. Nephrology Department, University Hospital Germans Trias i Pujol, Badalona, (Barcelona), Catalonia, Spain


b. International Renal Research Institute of Vicenza, Vicenza, Italy

c. Organization and Information Systems Department, University Hospital Germans Trias i Pujol, Badalona, (Barcelona), Catalonia, Spain



**Cr
Alert System
CrAS**

OBJECTIVE



**Cr
Alert System
CrAS**

METHOD



2019
2020
2021



Nephrology
Critical care
ER
CKDS5D

**SCr >1.5x baseline
SCr: >0.3mg/dL
Cr Alert System**

RESULTS






60,002
discharges

Discharges
23,180 (38.6%)
22,181 (36.9%)
23,641 (39.4%)

AKI per year
5,438 (9.1%)
AKI using CrAS

1,774 (32.6%)
1,778 (32.7%)
1. 886 (34.7%)



THE PAST

**Cr
Alert System**

**Easy and
Reliable
Tool**

THE FUTURE – AKI prevention

**Nephrology Rapid
Response Team**

**PoCUS
Biomarkers**



**40th on AKI&CRRT
VICENZA COURSE**
SPECIAL EDITION
June 14-16, 2022